

# Chapitre 1 : Introduction

Yassine ARIBA

$$\dot{x} = f(x)$$

# Sommaire

---

- ① Nonlinear models ?
- ② Existence of a solution
- ③ Equilibrium point
- ④ Linearization
- ⑤ Case study

# Sommaire

---

- 1 Nonlinear models ?
- 2 Existence of a solution
- 3 Equilibrium point
- 4 Linearization
- 5 Case study

## Linear systems

What you have seen so far... Models of the form

Linear ordinary differential equ.

$$\ddot{y}(t) + 3\dot{y}(t) + 2y(t) = u(t)$$

Transfer function

$$\frac{Y(s)}{U(s)} = \frac{1}{s^2 + 3s + 2}$$

Linear state space

$$\begin{aligned} \dot{x}(t) &= Ax(t) + Bu(t) \\ y(t) &= Cx(t) + Du(t) \end{aligned}$$


Fundamental property : superposition principle

$$\left\{ \begin{array}{l} u_1(t) \xrightarrow{\text{lin. sys.}} y_1(t) \\ u_2(t) \xrightarrow{\text{lin. sys.}} y_2(t) \end{array} \right. \Rightarrow a u_1(t) + b u_2(t) \xrightarrow{\text{lin. sys.}} a y_1(t) + b y_2(t)$$

↔ Allows to establish very strong and generic results

## Example

Consider a mass spring damper system



A simple model is obtained from Newton's law

$$m\ddot{y} + c\dot{y} + ky = 0$$

One can derive a Laplace domain or a state space representation

$$Y(s) = \frac{y_0}{ms^2 + cs + k}$$

$$\dot{x} = \begin{pmatrix} 0 & 1 \\ -\frac{k}{m} & -\frac{c}{m} \end{pmatrix} x$$

$$y = (1 \quad 0) x$$

## Nonlinear systems

More realistic models

$$\dot{x} = f(t, x, u)$$

where  $x$  is the state vector,  $u$  the input vector,  $f(\cdot)$  a **nonlinear** function.

Other cases :


- ▶ Unforced system :  $\dot{x} = f(t, x)$
- ▶ Autonomous system :  $\dot{x} = f(x)$  (case considered in the following)
- ▶ Affine in  $u$  :  $\dot{x} = f(x) + g(x)u$

Such a general modeling enables to better capture features of physical systems

↔ However, there is no general methods to deal with all nonlinear systems

## Example 1

Liquid level control



$S_1$  : section réservoir

$S_2$  : section sortie

$\rho$  : densité fluide

$P_1$  : pression en haut réservoir

$P_2$  : pression en sortie

$v_1$  : vitesse en haut réservoir

$v_2$  : vitesse en sortie

The change in mass in the tank is

$$\dot{m}(t) = \rho S_1 \dot{h}(t) = q_e(t) - \underbrace{\rho S_2 v_2(t)}_{q_s(t)}$$

Using the Bernouilli's equation :  $\frac{1}{2} \rho v_1^2(t) + P_1 + \rho g h(t) = \frac{1}{2} \rho v_2^2(t) + P_2$


$$\hookrightarrow v_2(t) = \sqrt{2gh(t)}$$

Let us define the state variable  $x = h$ , we get :

$$\dot{x}(t) = -a\sqrt{x(t)} + \frac{1}{\rho} q_e(t), \quad \text{with } a = \frac{S_2}{S_1} \sqrt{2g}$$

## Example 2

A simple free pendulum



Applying the Newton's second law, the equation of motion is obtained :

$$ml \ddot{\theta}(t) = -mg \sin \theta(t) - kl \dot{\theta}(t)$$


Let us define the state variables  $x_1 = \theta$  and  $x_2 = \dot{\theta}$ , we get :

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} x_2 \\ -\frac{g}{l} \sin x_1 - \frac{k}{m} x_2 \end{bmatrix}$$



## Origins of nonlinearities

- ▶ **Physical modeling.** Inherent to laws of Physics as in previous examples
- ▶ **Engineering design.** Inherent to how the system work, introduced by the engineer, technological aspect...



## Nonlinear phenomena


...that do not exist with linear modeling.

- ▶ **Multiple isolated equilibria.** Pendulum example
- ▶ **Finite escape time.** The state goes to infinity when time approaches a finite value. Example :

$$\dot{x} = -x^2, \quad \text{with the initial condition } x(0) = -1$$

⇒ The solution is

$$x(t) = \frac{1}{t-1}$$



## Nonlinear phenomena



### ► Limit cycles.

**Linear case** LTI systems oscillate if they have pure imaginary poles.

↪ It is a critical stability and nonrobust condition

↪ Oscillation amplitude depends on initial condition

$$\begin{cases} \dot{x}_1 = x_2 \\ \dot{x}_2 = -x_1 \end{cases}$$





**Nonlinear case** Can produce stable oscillations

↪ with fixed amplitude and frequency independently from initial conditions

Van der Pol equation

$$\begin{cases} \dot{x}_1 = x_2 \\ \dot{x}_2 = -x_1 + (1 - x_1^2)x_2 \end{cases}$$





## Nonlinear phenomena

### ► Frequency response

**Linear case** The response to a sine function is also a sine function (at steady state)



↪ with the same frequency  $\omega$

↪ and different amplitude and phase shift w.r.t.  $\omega$



**Nonlinear case** Can produce harmonics, subharmonics, and even almost-periodic output

$$\begin{cases} \dot{x}_1 = x_2^3 \\ \dot{x}_2 = -x_1 + (1 - x_2)u \end{cases}$$



# Sommaire

---

- ① Nonlinear models ?
- ② Existence of a solution
- ③ Equilibrium point
- ④ Linearization
- ⑤ Case study

## Existence of a solution

**Question** : (Cauchy problem)

Let be the system

$$\dot{x} = f(t, x), \quad \text{with the initial condition } x(t_0) = x_0 \in \mathbb{R}^n$$

Does a solution  $x(t)$  exist for  $t > t_0$ ? Is it unique? dependence on init. cond.?

### Theorem : local existence and uniqueness

If  $f(t, x)$  is piecewise continuous in  $t$  and satisfy the **Lipschitz condition**, that is, there exists a constant  $L > 0$  such that  $\forall x_1, x_2 \in B = \{x \in \mathbb{R}^n \mid \|x - x_0\| \leq r\}$ , and  $\forall t \in [t_0, t_1]$

$$\|f(t, x_2) - f(t, x_1)\| < L\|x_2 - x_1\|$$

then, there exists some  $\delta > 0$  such that the above system has a **unique solution** over  $[t_0, t_0 + \delta]$ .

## Example 1



$$\dot{x} = -x^3 \quad \text{with } x(0) = 1$$

$-x^3$  is Lipschitz for all  $x$  such that  $|x - x_0| \leq r = 1.5$

$$\frac{|-x_2^3 - (-x_1^3)|}{|x_2 - x_1|} \leq L$$

(but not true  $\forall x \in \mathbb{R}$ )

⇒ It exists a unique solution :  $x(t) = \frac{1}{\sqrt{1+2t}}$



## Example 2


$$\dot{x} = x^{2/3} \quad \text{with } x(0) = 0$$

has two solutions (non unicity) :  $x(t) = 0$  and  $x(t) = \frac{1}{27}t^3$ .

⇒ actually,  $x^{2/3}$  not Lipschitz around 0

$$\frac{|x^{2/3} - 0|}{|x - 0|} = |x^{-1/3}|$$

(not bounded when  $x \rightarrow 0$ )





## Example 3

$$\dot{x} = -x^2, \quad \text{with } x(0) = -1$$

$\Rightarrow -x^2$  is Lipschitz for  $\forall x_1, x_2 \in B = \{x \in \mathbb{R} \mid |x - x_0| \leq r\}$


$$\frac{|-x_2^2 - (-x_1^2)|}{|x_2 - x_1|} \leq L$$

(locally Lipschitz  $\forall x \in \mathbb{R}$ )

$\Rightarrow$  a unique solution for  $t \in [0, \delta]$


$$x(t) = \frac{1}{t - 1}$$

but  $\delta < 1$



## Lipschitz condition and derivative of $f$

Scalar and autonomous example :  $\dot{x} = f(x)$  with  $x \in \mathbb{R}$



A unique solution exists if

$$\frac{|f(x_2) - f(x_1)|}{|x_2 - x_1|} = \alpha \leq L \quad \forall x_1, x_2 \in B = \{x \in \mathbb{R} \mid |x - x_0| \leq r\}$$

$\hookrightarrow$  then  $f(x)$  is Lipschitz if  $|f'(x)|$  is bounded by  $L$

## Lipschitz condition and derivative of $f$

This observation extends to vector-valued functions

$$\left\| \frac{\partial f}{\partial x}(t, x) \right\| \leq L \quad f \text{ is Lipschitz} \quad (\text{for some domain})$$

### Lemma : Locally Lipschitz

If  $f(t, x)$  and  $\frac{\partial f}{\partial x}(t, x)$  are continuous on  $[t_0, t_1] \times D$ , for some domain  $D \subset \mathbb{R}^n$ , then  $f$  is locally Lipschitz on  $[t_0, t_1] \times D$ .

### Lemma : Globally Lipschitz

If  $f(t, x)$  and  $\frac{\partial f}{\partial x}(t, x)$  are continuous on  $[t_0, t_1] \times \mathbb{R}^n$ , then  $f$  is globally Lipschitz on  $[t_0, t_1] \times \mathbb{R}^n$  if and only if  $\frac{\partial f}{\partial x}$  is uniformly bounded on  $[t_0, t_1] \times \mathbb{R}^n$ .

## Back on previous examples

**Example 2 :**  $\dot{x} = x^{2/3}$ , with  $x(0) = 0$

$$\left(x^{2/3}\right)' = \frac{2}{3}x^{-1/3}$$

Hence,  $|f(x)'|$  unbounded at 0  $\Rightarrow f$  not Lipschitz around 0

**Example 3 :**  $\dot{x} = -x^2$ , with  $x(0) = -1$

$$\left(-x^2\right)' = -2x$$

Hence,  $|f(x)'|$  bounded for any  $x$  in some domain  $D \Rightarrow f$  locally Lipschitz  $\forall x \in \mathbb{R}$

## Exercise

Consider system

$$\dot{x} = f(x) = -x^2 + a \sin(x)$$

Is  $f(x)$  Lipschitz (locally or globally) or not ?

## Exercise

Consider system

$$\dot{x} = \underbrace{\begin{bmatrix} -x_1 + x_1x_2 \\ x_2 - x_1x_2 \end{bmatrix}}_{f(x)}$$

Is  $f(x)$  Lipschitz (locally or globally) or not ?

## Exercise

Consider system

$$\dot{x} = f(x) = -x + a \sin(x)$$

Is  $f(x)$  Lipschitz (locally or globally) or not ?

# Sommaire

---

- ① Nonlinear models ?
- ② Existence of a solution
- ③ Equilibrium point
- ④ Linearization
- ⑤ Case study



## Equilibrium point

### Definition

A point  $x^*$  is an **equilibrium point** if when the current state  $x = x^*$ , the system remains at this point ( $\rightarrow \dot{x} = 0$ ). The equilibrium points are given by the roots of

$$f(x) = 0$$

For the pendulum example, equilibrium points are characterized by

$$\begin{cases} 0 = x_2 \\ 0 = -\frac{g}{l} \sin x_1 - \frac{k}{m} x_2 \end{cases} \Rightarrow \begin{cases} x_2^* = 0 \\ x_1^* = 0 \pm n\pi, \quad n = 0, 1, 2, \dots \end{cases}$$

$\leftrightarrow$  mathematically infinitely many points, physically two positions

## Exercise

Consider system

$$\begin{cases} \dot{x}_1 = x_2 \\ \dot{x}_2 = -x_1(1 - a^2 x_1^2) - x_2 \end{cases}$$

where  $a > 0$  is a constant parameter.

Calculate the equilibrium point(s) ?

**Reminder** : for linear systems

$$\dot{x} = Ax + Bu \quad (A \text{ being non-singular})$$

there can be only one isolated equilibrium point  $x^* = -A^{-1}Bu^*$ .

- ▶ This equilibrium point is 0 in the case of an unforced system  $\dot{x} = Ax$ .
- ▶ If  $A$  is singular, there are infinitely many continuous equilibrium points (not isolated), this set is a subspace in the state-space.

# Sommaire

---

- ① Nonlinear models ?
- ② Existence of a solution
- ③ Equilibrium point
- ④ Linearization
- ⑤ Case study


## Linearization

Linear approximation of a nonlinear model around an equilibrium point

Pendulum example :

$$ml \ddot{\theta} = -mg \sin \theta - kl \dot{\theta}$$

around  $\theta = 0$ ,  $\sin(\theta) \sim \theta$



A **linear model** is obtained :

$$\begin{aligned}
 \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} &= \begin{bmatrix} x_2 \\ -\frac{g}{l} \sin x_1 - \frac{k}{m} x_2 \end{bmatrix} \xrightarrow{\text{around } \theta=0} \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} \simeq \begin{bmatrix} x_2 \\ -\frac{g}{l} x_1 - \frac{k}{m} x_2 \end{bmatrix} \\
 & \simeq \underbrace{\begin{bmatrix} 0 & 1 \\ -\frac{g}{l} & -\frac{k}{m} \end{bmatrix}}_A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}
 \end{aligned}$$

## More generally

Let's consider an equilibrium point  $x^*$  for system

$$\dot{x} = f(x), \quad \text{with } x(0) = x_0$$

and define the deviation variable :  $\tilde{x} = x - x^*$

Its dynamic is

$$\dot{\tilde{x}} = \dot{x} = f(x) = f(x^* + \tilde{x}), \quad \text{with } \tilde{x}(0) = x_0 - x^*$$


Use **Taylor series** around  $x^*$

$$f(x^* + h) = f(x^*) + f'(x^*)h + \frac{1}{2!}f''(x^*)h^2 + \frac{1}{3!}f^{(3)}(x^*)h^3 + \dots$$

valid if  $h (= \tilde{x})$  small enough

## Linear approximation (scalar case)

$$f(x^* + h) \simeq \underbrace{f(x^*)}_{=0} + f'(x^*)h + \frac{1}{2!} \cancel{f''(x^*)h^2} + \frac{1}{3!} \cancel{f^{(3)}(x^*)h^3} + \dots$$



For our system

$$\begin{aligned} \dot{\tilde{x}} &= f(x^* + \tilde{x}) \\ &\simeq f'(x^*) \tilde{x} \end{aligned}$$

⇒ **linear model** of the form :  $\dot{\tilde{x}} \simeq a \tilde{x}$ , with  $\tilde{x}(0) = x_0 - x^*$

## Linear approximation (general case)

Let  $x^*$  be an equilibrium point for system  $\dot{x} = f(x)$ , a linear model around that point is given by :

$$\dot{\tilde{x}} \simeq \underbrace{\frac{\partial f}{\partial x}(x^*)}_A \tilde{x} \quad \text{with } \tilde{x} = x - x^*$$

and  $\frac{\partial f}{\partial x}(\cdot)$  the Jacobian matrix of the vector-valued function  $f$  at the equ. pt.

- ▶ Reminder, Jacobian matrix :

$$\frac{\partial f}{\partial x}(x) = \begin{bmatrix} \frac{\partial f}{\partial x_1} & \dots & \frac{\partial f}{\partial x_n} \end{bmatrix} = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \dots & \frac{\partial f_1}{\partial x_n} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & & \vdots \\ \vdots & & \ddots & \vdots \\ \frac{\partial f_n}{\partial x_1} & \frac{\partial f_n}{\partial x_2} & \dots & \frac{\partial f_n}{\partial x_n} \end{bmatrix}$$

- ▶ One could also linearize around an operating point or a trajectory



## Back on the liquid level example


**Nonlinear model :**

$$\dot{x}(t) = -a\sqrt{x(t)} + \frac{1}{\rho}q_e(t), \quad \text{with } x(0) = 0.5 \text{ m}$$

For a constant input mass flow rate  $q_e(t) = q_0 \text{ kg/s} \Rightarrow$  equilibrium pt  $x^* = \left(\frac{q_0}{a\rho}\right)^2$

**Linearization :**

$$\dot{\tilde{x}} = -\frac{a}{2\sqrt{x^*}} \tilde{x}$$



## Back on the pendulum example

Nonlinear model :

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} x_2 \\ -\frac{g}{l} \sin x_1 - \frac{k}{m} x_2 \end{bmatrix}$$

Consider the equilibrium pt  $x^* = [0 \quad 0]^T$

Linearization :

Jacobian matrix

$$\frac{\partial f}{\partial x}(x) = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -\frac{g}{l} \cos x_1 & -\frac{k}{m} \end{bmatrix}$$

Linear model

$$\dot{\tilde{x}} = \begin{bmatrix} 0 & 1 \\ -\frac{g}{l} & -\frac{k}{m} \end{bmatrix} \tilde{x}$$

## Exercise

Consider system

$$\begin{cases} \dot{x}_1 = -x_1 + x_1 x_2 \\ \dot{x}_2 = x_1 + x_2 - 2x_1 x_2 \end{cases}$$

Calculate the equilibrium point(s)? Linearize the system around (1, 1)

# Sommaire

---

- ① Nonlinear models ?
- ② Existence of a solution
- ③ Equilibrium point
- ④ Linearization
- ⑤ Case study

## Case study

**Population dynamics** study the evolution of the size  $N(t)$  of a population

First simple model : *Malthus model*

$$\dot{N}(t) = \alpha N(t) - \beta N(t)$$

$\alpha$  is the birth rate and  $\beta$  the death rate

- ▶ Model is linear are nonlinear ?
- ▶ What is (are) the equilibrium point(s) ?
- ▶ Existence and unicity of the solution ?

## Solution

## Second case

Second model : *Verhulst (or logistic) model*

$$\dot{N}(t) = rN(t) \left( 1 - \frac{N(t)}{K} \right)$$

that takes into account a maximal critical size of the population  $K$  (carrying capacity).  $r$  is the growth rate.

- ▶ Model is linear or nonlinear ?
- ▶ What is (are) the equilibrium point(s) ?
- ▶ Existence and unicity of the solution ?

## Solution





## In short

- ▶ Nonlinear model are very general model

$$\dot{x} = f(x)$$

Results for linear model  $\dot{x} = Ax$  not applicable

- ▶ A solution exists and is unique if a Lipschitz condition is satisfied.
- ▶ The equilibrium points  $x^*$  are given by the roots of

$$f(x) = 0$$

- ▶ A nonlinear system may be approximated by a linear system around an equilibrium point

$$\dot{x} = f(x) \quad \xrightarrow{\text{approx}} \quad \delta\dot{x} = A \delta x$$

with  $x = x^* + \delta x$  and  $A = \frac{\partial f}{\partial x}(x^*)$